



Ardyan®

EMULSIFIABLE LIQUID

TECHNICAL DATASHEET

FICHA TÉCNICA

PERCENT COMPOSITION

% WEIGHT

ACTIVE INGREDIENT

Jasmonates (Cisjamone)

(equivalent to 6.5 g a.i./L at 20°C)

0.50

Chitosan (crustacean shell powder)

(equivalent to 26 g a.i./L at 20°C)

2.00

INERT INGREDIENTS

Conditioners, Stabilizers, Surfactant, Emulsifier, Adjuvant and Diluent.

97.50

TOTAL

100.00

RSCO-FUNG-0301H-0474-096-2.50

OMRI anm-18713

STOP, READ LABEL BEFORE USE PRODUCT MAY BE HARMFUL IF
SWALLOWED.

MAY BE HARMFUL IF IN DIRECT CONTACT WITH SKIN.

PREPARATION AND APPLICATION METHODS.

Shake the product well, carefully open the container, measure the indicated amount of Ardyan® with a plastic measuring cylinder or other suitable measuring instrument and pour it into a container half full with clean water, shake and then add it to the spraying tank or spray backpack and shake again.

Ardyan® is an emulsifiable liquid, useful for the control of the disease described below:

CROP	PEST	DOSE (L/ha)	NOTES
Cucumber Zucchini Watermelon Melon (0)	Powdery mildew (<i>Erysiphe cichoracearum</i>)	1.25 – 1.50	Apply 3 times to the foliage, at intervals of 7 and 11 days after the first application. Application volume 180-280 L of water/ha.

(): Safety interval: days from last application to harvest.

TIME OF RE-ENTRY INTO TREATED AREAS: Once the spray has dried.