

**Burhan®**

EMULSIFIABLE LIQUID

TECHNICAL DATASHEET

FICHA TÉCNICA

PERCENT COMPOSITION

% IN WEIGHT

ACTIVE INGREDIENTS

***Bacillus spp* (containing no less than 1×10^8 CFU/mL)**
(equivalent to 500 g a.i./L at 20°C)

50.00

Clove oil (*Syzygium aromaticum*)
(equivalent to 25 g a.i./L at 20°C)

2.50

Neem oil (*Azadirachta indica*)
(equivalent to 25 g a.i./L at 20°C)

2.50

INERT INGREDIENTS

Emulsifiers, Conditioners and Thinners

45.00

TOTAL

100.00

RSCO-MEZC-FUNG-1304-X0017-096-055

OMRI. anm-18819

STOP, READ LABEL BEFORE USE
MAY BE HARMFUL IF SWALLOWED, MAY BE HARMFUL IF IN CONTACT WITH SKIN.**ALWAYS CALIBRATE YOUR APPLICATION EQUIPMENT.**

Shake the product well; to open the container, twist the cap until the plastic ring is broken, then carefully remove the safety seal placed on top of the container. Measure with a plastic measuring cylinder or with some other suitable measuring instrument. Pour the specified quantity of BURHAN into a container containing half of its capacity of clean water, shake and then add it to the application tank or spraying backpack and shake again.

CROP	PEST	DOSE (L/ha)	NOTES
Coffee (0)	Coffee leaf rust (<i>Hemileia vastatrix</i> L.)	1.0-1.5	Apply four times to the foliage at a 14-day interval. Use the recommended dose in 1250-1350 L of water/ha

(0): Safety interval: days from last application to harvest.**TIME OF RE-ENTRY INTO TREATED AREAS:** Once the spray has dried.